arXiv:0709.0194v1 [math-ph] 3 Sep 2007

Gradings on $o(8,\mathbb{C})$

Cristina Draper *
Departamento de Matemática Aplicada, Universidad de Málaga
cdf@uma.es

Antonio Viruel†
Departamento de Álgebra, Geometría y Topología, Universidad de Málaga
viruel@agt.cie.uma.es

February 1, 2008

Abstract

This is a matricial description of all the fine group gradings on the exceptional Lie algebra $o(8,\mathbb{C})$. There are fourteen.

Keywords: fine grading, maximal abelian diagonalizable subgroup of automorphisms, exceptional Lie algebra, $\mathfrak{d}_4$.

1 Introduction

Given a Lie algebra L, a grading on L is a decomposition of vector spaces $L = \oplus_{i\in I} L_i$ such that $L_i \neq 0$ and for each pair of indices $i, j \in I$ there exists $k \in I$ such that $L_iL_j \subset L_k$ (see Definition 2.1 below). Besides of its own mathematical interest, gradings on Lie algebras show up in numerous applications of physical nature.

As it was already pointed out by Patera et al. in a series of papers on the subject [8, 9, 10, 11, 12], gradings are in the background of the choice of bases and the additive quantum numbers, and are a key ingredient when studying contractions that keep "undeformed" certain subalgebra (e.g. see [7]).

Gradings on Lie algebras also play a relevant role in quantum mechanics via Jordan algebras [6]: given a pair of indices $i, j \in I$ verifying $(L_iL_j)L_i \subset L_i$, any element $k \in L_j$ gives rise to a Jordan algebra structure in L_i by defining $x \circ y = (xk)y$.

Gradings on the algebras $o(n,\mathbb{C})$ have been computed in [1] and [5] for $n \neq 8$. The case $n = 8$ is of special relevance because of its strong symmetry, reflected partly by the triality automorphism. This makes that our matrices algebra is usually considered as a exceptional Lie algebra. Some works with a treatment of gradings on exceptional Lie algebras are [2, 3]. Here we describe in detail all the fourteen fine gradings on $o(8,\mathbb{C})$ (see Section 3). The proof showing that these are all the fine gradings is lengthy and cannot be included here, so it will appear in [4].

2 Basic concepts

In this minimal section, we introduce the basic concepts we use along this manuscript.

---

[*] Supported by MCYT grants MTM2004-06580-C02-02 and MTM2004-08115-C04-04, and by JA grants FQM-336 and FQM-1215

[†] Partially supported by the MCYT grant MTM2007-60016, and by the JA grant FQM-213

Definition 2.1. A *grading* on a Lie algebra L over a group G is a decomposition in vector spaces $L = \oplus_{g\in G} L_g$ such that $L_g L_h \subset L_{gh}$ for all $g, h \in G$ (sometimes called group grading). It will also be required that G is generated by $\{g \in G \mid L_g \neq 0\}$. We say that a grading is of *type* $(h_1, \dots, h_s)$ when there are h_i components of dimension i, and s is the greatest nonzero dimension of a component.

Definition 2.2. A grading is called *fine* when it cannot be refined. A *refinement* of the grading $L = \oplus_{g\in G} L_g$ is another (group) grading $L = \oplus_{h\in H} M_h$ such that each component L_g is sum of several components M_h.

Example 2.1. The main example of a fine grading is the root decomposition in a simple Lie algebra, which turns out to be a grading over $\mathbb{Z}^l$, for l the rank of the algebra.

From now on, L will denote the algebra of skewsymmetric matrices $o(8, \mathbb{C}) = \{x \in Mat_{8\times 8}(\mathbb{C}) \mid x + x^t = 0\}$. Let e_{ij} be the elementary matrix with (i, j)-entry 1, and the remaining entries zero, and define $b_{ij} = e_{ji} - e_{ij}$. Then $\{b_{ij} \mid i, j = 1, \dots 8, i < j\}$ is a basis of L, and the following section describes the fine gradings on L in terms of this base.

Recall that a grading on L is always the simultaneous diagonalization relative to a commuting set of semisimple automorphisms (more details in [2]). The grading is fine when it is produced by a maximal subgroup of semisimple automorphisms, which is usually called a *MAD*.

3 The fine gradings

We introduce some useful elements of $O(8, \mathbb{C})$ to describe the gradings on L.

$$\begin{array}{rl}
g(a) = & e_{11} + e_{22} + e_{33} + e_{44} + \frac{1}{2}(a + \frac{1}{a})(e_{55} + e_{66} + e_{77} + e_{88}) + \frac{i}{2}(a - \frac{1}{a})(e_{57} + e_{68} + e_{75} + e_{86}) \\
f(a) = & \frac{1}{2}(a + \frac{1}{a})(e_{11} + e_{22} + e_{33} + e_{44}) + (e_{55} + e_{66} + e_{77} + e_{88}) + \frac{i}{2}(a - \frac{1}{a})(e_{13} + e_{24} + e_{31} + e_{42}) \\
h(a) = & \frac{1}{2}(a + \frac{1}{a})(e_{11} + e_{22} + e_{33} + e_{44} + e_{55} + e_{66} + e_{77} + e_{88}) \\
 & +\frac{i}{2}(a - \frac{1}{a})(e_{15} + e_{26} + e_{37} + e_{48} + e_{51} + e_{62} + e_{73} + e_{84}) \\
p(a) = & e_{11} + e_{22} + e_{33} + e_{44} + e_{55} + e_{66} + \frac{1}{2}(a + \frac{1}{a})(e_{77} + e_{88}) + \frac{i}{2}(a - \frac{1}{a})(e_{78} + e_{87}) \\
q(a) = & e_{11} + e_{22} + e_{33} + e_{44} + \frac{1}{2}(a + \frac{1}{a})(e_{55} + e_{66}) + \frac{i}{2}(a - \frac{1}{a})(e_{56} + e_{65}) + e_{77} + e_{88} \\
r(a) = & e_{11} + e_{22} + \frac{1}{2}(a + \frac{1}{a})(e_{33} + e_{44}) + \frac{i}{2}(a - \frac{1}{a})(e_{34} + e_{43}) + e_{55} + e_{66} + e_{77} + e_{88} \\
s(a) = & \frac{1}{2}(a + \frac{1}{a})(e_{11} + e_{22}) + \frac{i}{2}(a - \frac{1}{a})(e_{12} + e_{21}) + e_{33} + e_{44} + e_{55} + e_{66} + e_{77} + e_{88}
\end{array}$$

$$\begin{array}{ll}
f_1 = \mathrm{diag}\{1, 1, 1, 1, 1, 1, 1, -1\} & f_2 = \mathrm{diag}\{1, 1, 1, 1, 1, 1, -1, 1\} \\
f_3 = \mathrm{diag}\{1, 1, 1, 1, 1, -1, 1, 1\} & f_4 = \mathrm{diag}\{1, 1, 1, 1, -1, 1, 1, 1\} \\
f_5 = \mathrm{diag}\{1, 1, 1, -1, 1, 1, 1, 1\} & f_6 = \mathrm{diag}\{1, 1, -1, 1, 1, 1, 1, 1\} \\
f_7 = \mathrm{diag}\{1, -1, 1, 1, 1, 1, 1, 1\} & f_8 = \mathrm{diag}\{-1, 1, 1, 1, 1, 1, 1, 1\}
\end{array}$$

$$\begin{array}{lllll}
g_1 = f_8 f_7 & g_2 = f_6 f_5 & g_4 = f_7 f_5 f_4 f_2 & g_5 = f_8 f_6 f_4 f_2 & \\
g_6 = f_2 f_1 & g_7 = f_7 f_5 f_3 f_1 & g_9 = f_8 f_7 f_4 f_3 & g_{10} = f_6 f_5 f_2 f_1 & g_{13} = f_4 f_3 f_2 f_1
\end{array}$$

$$\begin{array}{rl}
g_3 = & e_{12} + e_{21} + e_{34} + e_{43} + e_{56} + e_{65} + e_{78} + e_{87} \\
g_8 = & e_{13} + e_{31} + e_{24} + e_{42} + e_{57} + e_{75} + e_{68} + e_{86} \\
g_{11} = & e_{12} + e_{21} + e_{34} + e_{43} - e_{55} + e_{66} - e_{77} + e_{88} \\
g_{12} = & e_{11} - e_{22} + e_{33} - e_{44} - e_{56} + e_{65} - e_{78} + e_{87} \\
g_{14} = & e_{15} + e_{26} + e_{37} + e_{48} + e_{51} + e_{62} + e_{73} + e_{84}
\end{array}$$

Notice that $f_i, g_i, f(a), g(a), h(a), p(a), q(a), r(a), s(a) \in O(8, \mathbb{C}) = \{P \in Mat_{8\times 8}(\mathbb{C}) \mid PP^t = \mathrm{id}\}$ for all $a \in \mathbb{C}^*$. Thus we can use the adjoint map to obtain automorphisms of L by means of $\mathrm{Ad}\colon O(8, \mathbb{C}) \to \mathrm{Aut}\, o(8, \mathbb{C})$, $\mathrm{Ad}\, P(x) = P^{-1} x P$. In particular, denote $F_i = \mathrm{Ad}\, f_i$ and $G_i = \mathrm{Ad}\, g_i$. With all these elements, we are prepared to describe completely the group of automorphisms producing each grading, denoted by Q_i. We also provide the simultaneous diagonalization, which is a straightforward computation by means of any computer algebra software.

Grading over $\mathbb{Z}\times\mathbb{Z}_2^4$: $Q_1 = \langle\{\operatorname{Ad} g(u), G_1, G_2, G_3, G_4 \mid u \in \mathbb{C}^*\}\rangle$ is an abelian diagonalizable subgroup of $\operatorname{Aut} L$. In fact it is maximal according to [5] (it is the MAD corresponding to $T_{2,2}^{(1)}$). Hence it produces a fine grading, of type $(25, 0, 1)$, whose homogeneous components are:

$$\begin{array}{ll}
L_{\frac{1}{2},1,-1,-1,-1} = \langle ib_{3,5} - b_{3,7} - ib_{4,6} + b_{4,8}\}\rangle & L_{\frac{1}{4},1,1,-1,-1} = \langle -b_{5,6} - ib_{5,8} + ib_{6,7} + b_{7,8}\rangle \\
L_{\frac{1}{2},1,-1,-1,1} = \langle ib_{3,6} - b_{3,8} - ib_{4,5} + b_{4,7}\rangle & L_{\frac{1}{2},-1,1,-1,-1} = \langle ib_{1,5} - b_{1,7} - ib_{2,6} + b_{2,8}\rangle \\
L_{\frac{1}{2},1,-1,1,-1} = \langle -ib_{3,5} + b_{3,7} - ib_{4,6} + b_{4,8}\rangle & L_{\frac{1}{2},-1,1,-1,1} = \langle ib_{1,6} - b_{1,8} - ib_{2,5} + b_{2,7}\rangle \\
L_{\frac{1}{2},1,-1,1,1} = \langle -ib_{3,6} + b_{3,8} - ib_{4,5} + b_{4,7}\rangle & L_{\frac{1}{2},-1,1,1,-1} = \langle -ib_{1,5} + b_{1,7} - ib_{2,6} + b_{2,8}\rangle \\
L_{\frac{1}{2},-1,1,1,1} = \langle -ib_{1,6} + b_{1,8} - ib_{2,5} + b_{2,7}\rangle & L_{1,1,1,-1,-1} = \langle b_{1,2},\, b_{3,4},\, b_{7,8} + b_{5,6}\rangle \\
L_{2,1,-1,-1,1} = \langle -ib_{3,6} - b_{3,8} + ib_{4,5} + b_{4,7}\rangle & L_{2,-1,1,-1,-1} = \langle -ib_{1,5} - b_{1,7} + ib_{2,6} + b_{2,8}\rangle \\
L_{2,1,-1,1,-1} = \langle ib_{3,5} + b_{3,7} + ib_{4,6} + b_{4,8}\rangle & L_{2,-1,1,-1,1} = \langle -ib_{1,6} - b_{1,8} + ib_{2,5} + b_{2,7}\rangle \\
L_{2,1,-1,1,1} = \langle ib_{3,6} + b_{3,8} + ib_{4,5} + b_{4,7}\rangle & L_{2,-1,1,1,-1} = \langle ib_{1,5} + b_{1,7} + ib_{2,6} + b_{2,8}\rangle \\
L_{4,1,1,-1,-1} = \langle -b_{5,6} + ib_{5,8} - ib_{6,7} + b_{7,8}\rangle & L_{2,-1,1,1,1} = \langle ib_{1,6} + b_{1,8} + ib_{2,5} + b_{2,7}\rangle \\
L_{1,1,1,1,-1} = \langle b_{5,8} + b_{6,7}\rangle & L_{2,1,-1,-1,-1} = \langle -ib_{3,5} - b_{3,7} + ib_{4,6} + b_{4,8}\rangle
\end{array}$$

$$\begin{array}{lll}
L_{1,-1,-1,-1,-1} = \langle b_{2,3} - b_{1,4}\rangle & L_{1,1,1,1,1} = \langle b_{6,8} + b_{5,7}\rangle & L_{1,-1,-1,1,-1} = \langle b_{2,3} + b_{1,4}\rangle \\
L_{1,-1,-1,-1,1} = \langle b_{2,4} - b_{1,3}\rangle & L_{1,1,1,-1,1} = \langle b_{6,8} - b_{5,7}\rangle & L_{1,-1,-1,1,1} = \langle b_{2,4} + b_{1,3}\rangle
\end{array}$$

The indices of the components are not the indices of the grading, but the eigenvalues that $\{\operatorname{Ad} g(2), G_1, G_2, G_3, G_4\}$ act with (e.g. the index $\{\frac{1}{2}, 1, -1, -1, -1\}$ is for the element $(-2, \bar{0}, \bar{1}, \bar{1}, \bar{1}) \in \mathbb{Z}\times\mathbb{Z}_2^4$). The universal group of the grading (see [2] for the definition) is just $\mathbb{Z}\times\mathbb{Z}_2^4 (= \chi(Q_1))$. These comments about the universal group and the order and eigenvalues of the automorphisms, are valid for all the gradings in this paper.

Grading over $\mathbb{Z}^2\times\mathbb{Z}_2^2$: $Q_2 = \langle\{\operatorname{Ad} f(u), \operatorname{Ad} g(v), G_3, G_5 \mid u, v \in \mathbb{C}^*\}\rangle \cong (\mathbb{C}^*)^2 \times \mathbb{Z}_2^2$ is an abelian diagonalizable subgroup of $\operatorname{Aut} L$. In fact, it is the MAD corresponding to $T_{4,0}^{(1)}$ ([5]). The simultaneous diagonalization of L relative to this subgroup of automorphisms is of type $(20, 4)$:

$$\begin{array}{l}
L_{\frac{1}{2},\frac{1}{3},-1,-1} = \langle b_{1,6} + ib_{1,8} - b_{2,5} - ib_{2,7} + ib_{3,6} - b_{3,8} - ib_{4,5} + b_{4,7}\rangle \\
L_{\frac{1}{2},\frac{1}{3},-1,1} = \langle b_{1,5} + ib_{1,7} - b_{2,6} - ib_{2,8} + ib_{3,5} - b_{3,7} - ib_{4,6} + b_{4,8}\rangle \\
L_{\frac{1}{2},\frac{1}{3},1,-1} = \langle -b_{1,6} - ib_{1,8} - b_{2,5} - ib_{2,7} - ib_{3,6} + b_{3,8} - ib_{4,5} + b_{4,7}\rangle
\end{array}$$

$$\begin{array}{l}
L_{\frac{1}{2},\frac{1}{3},1,1} = \langle -b_{1,5} - ib_{1,7} - b_{2,6} - ib_{2,8} - ib_{3,5} + b_{3,7} - ib_{4,6} + b_{4,8}\rangle \\
L_{\frac{1}{2},3,-1,-1} = \langle -b_{1,6} + ib_{1,8} + b_{2,5} - ib_{2,7} - ib_{3,6} - b_{3,8} + ib_{4,5} + b_{4,7}\rangle \\
L_{\frac{1}{2},3,-1,1} = \langle -b_{1,5} + ib_{1,7} + b_{2,6} - ib_{2,8} - ib_{3,5} - b_{3,7} + ib_{4,6} + b_{4,8}\rangle \\
L_{\frac{1}{2},3,1,-1} = \langle b_{1,6} - ib_{1,8} + b_{2,5} - ib_{2,7} + ib_{3,6} + b_{3,8} + ib_{4,5} + b_{4,7}\rangle \\
L_{\frac{1}{2},3,1,1} = \langle b_{1,5} - ib_{1,7} + b_{2,6} - ib_{2,8} + ib_{3,5} + b_{3,7} + ib_{4,6} + b_{4,8}\rangle \\
L_{2,\frac{1}{3},-1,-1} = \langle -b_{1,6} - ib_{1,8} + b_{2,5} + ib_{2,7} + ib_{3,6} - b_{3,8} - ib_{4,5} + b_{4,7}\rangle \\
L_{2,\frac{1}{3},-1,1} = \langle -b_{1,5} - ib_{1,7} + b_{2,6} + ib_{2,8} + ib_{3,5} - b_{3,7} - ib_{4,6} + b_{4,8}\rangle \\
L_{2,\frac{1}{3},1,-1} = \langle b_{1,6} + ib_{1,8} + b_{2,5} + ib_{2,7} - ib_{3,6} + b_{3,8} - ib_{4,5} + b_{4,7}\rangle \\
L_{2,\frac{1}{3},1,1} = \langle b_{1,5} + ib_{1,7} + b_{2,6} + ib_{2,8} - ib_{3,5} + b_{3,7} - ib_{4,6} + b_{4,8}\rangle \\
L_{2,3,-1,-1} = \langle b_{1,6} - ib_{1,8} - b_{2,5} + ib_{2,7} - ib_{3,6} - b_{3,8} + ib_{4,5} + b_{4,7}\rangle \\
L_{2,3,-1,1} = \langle b_{1,5} - ib_{1,7} - b_{2,6} + ib_{2,8} - ib_{3,5} - b_{3,7} + ib_{4,6} + b_{4,8}\rangle \\
L_{2,3,1,-1} = \langle -b_{1,6} + ib_{1,8} - b_{2,5} + ib_{2,7} + ib_{3,6} + b_{3,8} + ib_{4,5} + b_{4,7}\rangle \\
L_{2,3,1,1} = \langle -b_{1,5} + ib_{1,7} - b_{2,6} + ib_{2,8} + ib_{3,5} + b_{3,7} + ib_{4,6} + b_{4,8}\rangle
\end{array}$$

$$\begin{array}{ll}
L_{1,1,1,1} = \langle b_{1,3} + b_{2,4},\, b_{5,7} + b_{6,8}\rangle & L_{1,1,-1,-1} = \langle b_{1,2} + b_{3,4},\, b_{5,6} + b_{7,8}\rangle \\
L_{1,1,-1,1} = \langle -b_{1,3} + b_{2,4},\, -b_{5,7} + b_{6,8}\rangle & L_{1,1,1,-1} = \langle b_{1,4} + b_{2,3},\, b_{5,8} + b_{6,7}\rangle \\
L_{\frac{1}{4},1,-1,-1} = \langle -b_{1,2} - ib_{1,4} + ib_{2,3} + b_{3,4}\rangle & L_{1,9,-1,-1} = \langle -b_{5,6} + ib_{5,8} - ib_{6,7} + b_{7,8}\rangle \\
L_{4,1,-1,-1} = \langle -b_{1,2} + ib_{1,4} - ib_{2,3} + b_{3,4}\rangle & L_{1,\frac{1}{9},-1,-1} = \langle -b_{5,6} - ib_{5,8} + ib_{6,7} + b_{7,8}\rangle
\end{array}$$

Grading over $\mathbb{Z}_2^5$: $Q_3 = \langle\{G_6, G_1, G_2, G_3, G_7\}\rangle \cong \mathbb{Z}_2^5$ is a MAD of $\operatorname{Aut} L$, corresponding to $T_{0,4}^{(1)}$. The simultaneous diagonalization of L relative to the quasitorus Q_3 is of type $(24, 0, 0, 1)$, with $L_{1,1,1,-1,-1} = \langle b_{1,2}, b_{3,4}, b_{5,6}, b_{7,8}\rangle$ and:

$$\begin{array}{lll}
L_{-1,-1,1,-1,-1} = \langle -b_{1,8} + b_{2,7}\rangle & L_{-1,-1,1,-1,1} = \langle -b_{1,7} + b_{2,8}\rangle & L_{-1,-1,1,1,-1} = \langle b_{1,8} + b_{2,7}\rangle \\
L_{-1,-1,1,1,1} = \langle b_{1,7} + b_{2,8}\rangle & L_{-1,1,-1,-1,-1} = \langle -b_{3,8} + b_{4,7}\rangle & L_{-1,1,-1,-1,1} = \langle -b_{3,7} + b_{4,8}\rangle \\
L_{-1,1,-1,1,-1} = \langle b_{3,8} + b_{4,7}\rangle & L_{-1,1,-1,1,1} = \langle b_{3,7} + b_{4,8}\rangle & L_{-1,1,1,-1,-1} = \langle -b_{5,8} + b_{6,7}\rangle \\
L_{-1,1,1,-1,1} = \langle -b_{5,7} + b_{6,8}\rangle & L_{-1,1,1,1,-1} = \langle b_{5,8} + b_{6,7}\rangle & L_{-1,1,1,1,1} = \langle b_{5,7} + b_{6,8}\rangle \\
L_{1,-1,-1,-1,-1} = \langle -b_{1,4} + b_{2,3}\rangle & L_{1,-1,-1,-1,1} = \langle -b_{1,3} + b_{2,4}\rangle & L_{1,-1,-1,1,-1} = \langle b_{1,4} + b_{2,3}\rangle \\
L_{1,-1,-1,1,1} = \langle b_{1,3} + b_{2,4}\rangle & L_{1,-1,1,-1,-1} = \langle -b_{1,6} + b_{2,5}\rangle & L_{1,-1,1,-1,1} = \langle -b_{1,5} + b_{2,6}\rangle \\
L_{1,-1,1,1,-1} = \langle b_{1,6} + b_{2,5}\rangle & L_{1,-1,1,1,1} = \langle b_{1,5} + b_{2,6}\rangle & L_{1,1,-1,-1,-1} = \langle -b_{3,6} + b_{4,5}\rangle \\
L_{1,1,-1,-1,1} = \langle -b_{3,5} + b_{4,6}\rangle & L_{1,1,-1,1,-1} = \langle b_{3,6} + b_{4,5}\rangle & L_{1,1,-1,1,1} = \langle b_{3,5} + b_{4,6}\rangle
\end{array}$$

Grading over $\mathbb{Z} \times \mathbb{Z}_2^4$: $Q_4 = \langle\{\operatorname{Ad} h(u), G_8, G_9, G_3, G_5 \mid u \in \mathbb{C}^*\}\rangle \cong (\mathbb{C}^*) \times \mathbb{Z}_2^4$ is a MAD of $\operatorname{Aut} L$, corresponding to $T_{2,0}^{(2)}$. The simultaneous diagonalization of L relative to the quasitorus Q_4 is of type (28):

$$\begin{array}{ll}
L_{1,-1,-1,-1,-1} = \langle b_{1,8} - b_{2,7} - b_{3,6} + b_{4,5}\rangle & L_{1,-1,-1,-1,1} = \langle -b_{1,3} + b_{2,4} - b_{5,7} + b_{6,8}\rangle \\
L_{1,-1,-1,1,-1} = \langle b_{1,4} + b_{2,3} + b_{5,8} + b_{6,7}\rangle & L_{1,-1,-1,1,1} = \langle b_{1,3} + b_{2,4} + b_{5,7} + b_{6,8}\rangle \\
L_{1,-1,1,-1,-1} = \langle -b_{1,2} + b_{3,4} - b_{5,6} + b_{7,8}\rangle & L_{1,-1,1,-1,1} = \langle b_{1,5} - b_{2,6} - b_{3,7} + b_{4,8}\rangle \\
L_{1,-1,1,1,-1} = \langle -b_{1,6} - b_{2,5} + b_{3,8} + b_{4,7}\rangle & L_{1,-1,1,1,1} = \langle -b_{1,5} - b_{2,6} + b_{3,7} + b_{4,8}\rangle \\
L_{1,1,-1,-1,-1} = \langle -b_{1,4} + b_{2,3} - b_{5,8} + b_{6,7}\rangle & L_{1,1,-1,-1,1} = \langle -b_{1,7} + b_{2,8} - b_{3,5} + b_{4,6}\rangle \\
L_{1,1,-1,1,-1} = \langle b_{1,8} + b_{2,7} + b_{3,6} + b_{4,5}\rangle & L_{1,1,-1,1,1} = \langle b_{1,7} + b_{2,8} + b_{3,5} + b_{4,6}\rangle \\
L_{1,1,1,-1,-1} = \langle b_{1,2} + b_{3,4} + b_{5,6} + b_{7,8}\rangle & L_{1,1,1,-1,1} = \langle -b_{1,5} + b_{2,6} - b_{3,7} + b_{4,8}\rangle \\
L_{1,1,1,1,-1} = \langle b_{1,6} + b_{2,5} + b_{3,8} + b_{4,7}\rangle & L_{1,1,1,1,1} = \langle b_{1,5} + b_{2,6} + b_{3,7} + b_{4,8}\rangle
\end{array}$$

$$\begin{array}{l}
L_{\frac{1}{4},-1,-1,-1,1} = \langle b_{1,3} + ib_{1,7} - b_{2,4} - ib_{2,8} - ib_{3,5} + ib_{4,6} - b_{5,7} + b_{6,8}\rangle \\
L_{\frac{1}{4},-1,-1,1,-1} = \langle -b_{1,4} - ib_{1,8} - b_{2,3} - ib_{2,7} + ib_{3,6} + ib_{4,5} + b_{5,8} + b_{6,7}\rangle \\
L_{\frac{1}{4},-1,-1,1,1} = \langle -b_{1,3} - ib_{1,7} - b_{2,4} - ib_{2,8} + ib_{3,5} + ib_{4,6} + b_{5,7} + b_{6,8}\rangle \\
L_{\frac{1}{4},-1,1,-1,-1} = \langle b_{1,2} + ib_{1,6} - ib_{2,5} - b_{3,4} - ib_{3,8} + ib_{4,7} - b_{5,6} + b_{7,8}\rangle \\
L_{\frac{1}{4},1,-1,-1,-1} = \langle b_{1,4} + ib_{1,8} - b_{2,3} - ib_{2,7} + ib_{3,6} - ib_{4,5} - b_{5,8} + b_{6,7}\rangle \\
L_{\frac{1}{4},1,1,-1,-1} = \langle -b_{1,2} - ib_{1,6} + ib_{2,5} - b_{3,4} - ib_{3,8} + ib_{4,7} + b_{5,6} + b_{7,8}\rangle \\
L_{4,-1,-1,-1,1} = \langle b_{1,3} - ib_{1,7} - b_{2,4} + ib_{2,8} + ib_{3,5} - ib_{4,6} - b_{5,7} + b_{6,8}\rangle \\
L_{4,-1,-1,1,-1} = \langle -b_{1,4} + ib_{1,8} - b_{2,3} + ib_{2,7} - ib_{3,6} - ib_{4,5} + b_{5,8} + b_{6,7}\rangle \\
L_{4,-1,-1,1,1} = \langle -b_{1,3} + ib_{1,7} - b_{2,4} + ib_{2,8} - ib_{3,5} - ib_{4,6} + b_{5,7} + b_{6,8}\rangle \\
L_{4,-1,1,-1,-1} = \langle b_{1,2} - ib_{1,6} + ib_{2,5} - b_{3,4} + ib_{3,8} - ib_{4,7} - b_{5,6} + b_{7,8}\rangle \\
L_{4,1,-1,-1,-1} = \langle b_{1,4} - ib_{1,8} - b_{2,3} + ib_{2,7} - ib_{3,6} + ib_{4,5} - b_{5,8} + b_{6,7}\rangle \\
L_{4,1,1,-1,-1} = \langle -b_{1,2} + ib_{1,6} - ib_{2,5} - b_{3,4} + ib_{3,8} - ib_{4,7} + b_{5,6} + b_{7,8}\rangle
\end{array}$$

Grading over $\mathbb{Z}_2^7$: $Q_5 = \langle\{F_i \mid i = 1, \ldots, 7\}\rangle \cong \mathbb{Z}_2^7$ is a MAD of $\operatorname{Aut} L$, corresponding to $T_{0,8}^{(0)}$. The simultaneous diagonalization of L relative to it is of type (28):

$$\begin{array}{llll}
L_{1,-1,-1,1,1,1,1} = \langle b_{6,7}\rangle & L_{1,-1,1,-1,1,1,1} = \langle b_{5,7}\rangle & L_{1,-1,1,1,-1,1,1} = \langle b_{4,7}\rangle & L_{1,-1,1,1,1,-1,1} = \langle b_{3,7}\rangle \\
L_{1,-1,1,1,1,1,-1} = \langle b_{2,7}\rangle & L_{1,-1,1,1,1,1,1} = \langle b_{1,7}\rangle & L_{1,1,-1,-1,1,1,1} = \langle b_{5,6}\rangle & L_{1,1,-1,1,-1,1,1} = \langle b_{4,6}\rangle \\
L_{1,1,-1,1,1,-1,1} = \langle b_{3,6}\rangle & L_{1,1,-1,1,1,1,-1} = \langle b_{2,6}\rangle & L_{1,1,-1,1,1,1,1} = \langle b_{1,6}\rangle & L_{1,1,1,-1,-1,1,1} = \langle b_{4,5}\rangle \\
L_{1,1,1,-1,1,-1,1} = \langle b_{3,5}\rangle & L_{1,1,1,-1,1,1,-1} = \langle b_{2,5}\rangle & L_{1,1,1,-1,1,1,1} = \langle b_{1,5}\rangle & L_{1,1,1,1,-1,-1,1} = \langle b_{3,4}\rangle \\
L_{1,1,1,1,-1,1,-1} = \langle b_{2,4}\rangle & L_{1,1,1,1,-1,1,1} = \langle b_{1,4}\rangle & L_{1,1,1,1,1,-1,-1} = \langle b_{2,3}\rangle & L_{1,1,1,1,1,-1,1} = \langle b_{1,3}\rangle \\
L_{1,1,1,1,1,1,-1} = \langle b_{1,2}\rangle & L_{-1,-1,1,1,1,1,1} = \langle b_{7,8}\rangle & L_{-1,1,-1,1,1,1,1} = \langle b_{6,8}\rangle & L_{-1,1,1,-1,1,1,1} = \langle b_{5,8}\rangle \\
L_{-1,1,1,1,-1,1,1} = \langle b_{4,8}\rangle & L_{-1,1,1,1,1,-1,1} = \langle b_{3,8}\rangle & L_{-1,1,1,1,1,1,-1} = \langle b_{2,8}\rangle & L_{-1,1,1,1,1,1,1} = \langle b_{1,8}\rangle
\end{array}$$

Grading over $\mathbb{Z} \times \mathbb{Z}_2^5$: $Q_6 = \langle\{p(u), F_8, F_7, F_6, F_5, F_4 \mid u \in \mathbb{C}^*\}\rangle \cong \mathbb{C}^* \times \mathbb{Z}_2^5$ is again an abelian diagonalizable subgroup of $\operatorname{Aut} L$, concretely the MAD $T_{2,6}^{(0)}$. The simultaneous diagonalization of L

relative to the quasitorus Q_6 is of type (28):

$$\begin{array}{ll}
L_{\frac{1}{2},-1,1,1,1,1} = \langle ib_{1,7} + b_{1,8} \rangle & L_{\frac{1}{2},1,-1,1,1,1} = \langle ib_{2,7} + b_{2,8} \rangle \\
L_{\frac{1}{2},1,1,-1,1,1} = \langle ib_{3,7} + b_{3,8} \rangle & L_{\frac{1}{2},1,1,1,-1,1} = \langle ib_{4,7} + b_{4,8} \rangle \\
L_{\frac{1}{2},1,1,1,1,-1} = \langle ib_{5,7} + b_{5,8} \rangle & L_{\frac{1}{2},1,1,1,1,1} = \langle ib_{6,7} + b_{6,8} \rangle \\
L_{2,-1,1,1,1,1} = \langle -ib_{1,7} + b_{1,8} \rangle & L_{2,1,-1,1,1,1} = \langle -ib_{2,7} + b_{2,8} \rangle \\
L_{2,1,1,-1,1,1} = \langle -ib_{3,7} + b_{3,8} \rangle & L_{2,1,1,1,-1,1} = \langle -ib_{4,7} + b_{4,8} \rangle \\
L_{2,1,1,1,1,-1} = \langle -ib_{5,7} + b_{5,8} \rangle & L_{2,1,1,1,1,1} = \langle -ib_{6,7} + b_{6,8} \rangle
\end{array}$$

$$\begin{array}{llll}
L_{1,-1,-1,1,1,1} = \langle b_{1,2} \rangle & L_{1,-1,1,-1,1,1} = \langle b_{1,3} \rangle & L_{1,-1,1,1,-1,1} = \langle b_{1,4} \rangle & L_{1,-1,1,1,1,-1} = \langle b_{1,5} \rangle \\
L_{1,-1,1,1,1,1} = \langle b_{1,6} \rangle & L_{1,1,-1,-1,1,1} = \langle b_{2,3} \rangle & L_{1,1,-1,1,-1,1} = \langle b_{2,4} \rangle & L_{1,1,-1,1,1,-1} = \langle b_{2,5} \rangle \\
L_{1,1,-1,1,1,1} = \langle b_{2,6} \rangle & L_{1,1,1,-1,-1,1} = \langle b_{3,4} \rangle & L_{1,1,1,-1,1,-1} = \langle b_{3,5} \rangle & L_{1,1,1,-1,1,1} = \langle b_{3,6} \rangle \\
L_{1,1,1,1,-1,-1} = \langle b_{4,5} \rangle & L_{1,1,1,1,-1,1} = \langle b_{4,6} \rangle & L_{1,1,1,1,1,-1} = \langle b_{5,6} \rangle & L_{1,1,1,1,1,1} = \langle b_{7,8} \rangle
\end{array}$$

Grading over $\mathbb{Z}^2 \times \mathbb{Z}_2^3$: $Q_7 = \langle \{p(u), F_8, F_7, F_6, q(v) \mid u, v \in \mathbb{C}^*\} \rangle \cong (\mathbb{C}^*)^2 \times \mathbb{Z}_2^3$ is a MAD of $\operatorname{Aut} L$, corresponding to $T_{4,4}^{(0)}$, producing the following grading, of type $(26, 1)$:

$$\begin{array}{ll}
L_{2,1,1,1,3} = \langle -b_{5,7} - ib_{5,8} - ib_{6,7} + b_{6,8} \rangle & L_{\frac{1}{2},1,1,1,\frac{1}{3}} = \langle -b_{5,7} + ib_{5,8} + ib_{6,7} + b_{6,8} \rangle \\
L_{2,1,1,1,\frac{1}{3}} = \langle b_{5,7} + ib_{5,8} - ib_{6,7} + b_{6,8} \rangle & L_{\frac{1}{2},1,1,1,3} = \langle b_{5,7} - ib_{5,8} + ib_{6,7} + b_{6,8} \rangle
\end{array}$$

$$\begin{array}{lll}
L_{\frac{1}{2},-1,1,1,1} = \langle ib_{1,7} + b_{1,8} \rangle & L_{\frac{1}{2},1,-1,1,1} = \langle ib_{2,7} + b_{2,8} \rangle & L_{\frac{1}{2},1,1,-1,1} = \langle ib_{3,7} + b_{3,8} \rangle \\
L_{\frac{1}{2},1,1,1,1} = \langle ib_{4,7} + b_{4,8} \rangle & L_{1,-1,-1,1,1} = \langle b_{1,2} \rangle & L_{1,-1,1,-1,1} = \langle b_{1,3} \rangle \\
L_{1,-1,1,1,\frac{1}{3}} = \langle ib_{1,5} + b_{1,6} \rangle & L_{1,-1,1,1,1} = \langle b_{1,4} \rangle & L_{1,-1,1,1,3} = \langle -ib_{1,5} + b_{1,6} \rangle \\
L_{1,1,-1,-1,1} = \langle b_{2,3} \rangle & L_{1,1,-1,1,\frac{1}{3}} = \langle ib_{2,5} + b_{2,6} \rangle & L_{1,1,-1,1,1} = \langle b_{2,4} \rangle \\
L_{1,1,-1,1,3} = \langle -ib_{2,5} + b_{2,6} \rangle & L_{1,1,1,-1,\frac{1}{3}} = \langle ib_{3,5} + b_{3,6} \rangle & L_{1,1,1,-1,1} = \langle b_{3,4} \rangle \\
L_{1,1,1,-1,3} = \langle -ib_{3,5} + b_{3,6} \rangle & L_{1,1,1,1,\frac{1}{3}} = \langle ib_{4,5} + b_{4,6} \rangle & L_{1,1,1,1,1} = \langle b_{5,6},\ b_{7,8} \rangle \\
L_{1,1,1,1,3} = \langle -ib_{4,5} + b_{4,6} \rangle & L_{2,-1,1,1,1} = \langle -ib_{1,7} + b_{1,8} \rangle & L_{2,1,-1,1,1} = \langle -ib_{2,7} + b_{2,8} \rangle \\
L_{2,1,1,-1,1} = \langle -ib_{3,7} + b_{3,8} \rangle & L_{2,1,1,1,1} = \langle -ib_{4,7} + b_{4,8} \rangle &
\end{array}$$

Grading over $\mathbb{Z}_2^3 \times \mathbb{Z}_4$: $Q_8 = \langle \{G_8, G_{10}, G_{11}, G_{12}\} \rangle \cong \mathbb{Z}_2^3 \times \mathbb{Z}_4$ is a MAD of $\operatorname{Aut} L$, corresponding to $T_{0,2}^{(2)}$, which produces in L a grading of type $(24, 2)$:

$$\begin{array}{l}
L_{-1,-1,-1,-i} = \langle ib_{1,7} + b_{1,8} + ib_{2,7} - b_{2,8} - ib_{3,5} - b_{3,6} - ib_{4,5} + b_{4,6} \rangle \\
L_{-1,-1,-1,i} = \langle -ib_{1,7} + b_{1,8} - ib_{2,7} - b_{2,8} + ib_{3,5} - b_{3,6} + ib_{4,5} + b_{4,6} \rangle \\
L_{-1,-1,1,-i} = \langle -ib_{1,7} - b_{1,8} + ib_{2,7} - b_{2,8} + ib_{3,5} + b_{3,6} - ib_{4,5} + b_{4,6} \rangle \\
L_{-1,-1,1,i} = \langle ib_{1,7} - b_{1,8} - ib_{2,7} - b_{2,8} - ib_{3,5} + b_{3,6} + ib_{4,5} + b_{4,6} \rangle \\
L_{-1,1,-1,-i} = \langle ib_{1,5} + b_{1,6} + ib_{2,5} - b_{2,6} - ib_{3,7} - b_{3,8} - ib_{4,7} + b_{4,8} \rangle \\
L_{-1,1,-1,i} = \langle -ib_{1,5} + b_{1,6} - ib_{2,5} - b_{2,6} + ib_{3,7} - b_{3,8} + ib_{4,7} + b_{4,8} \rangle \\
L_{-1,1,1,-i} = \langle -ib_{1,5} - b_{1,6} + ib_{2,5} - b_{2,6} + ib_{3,7} + b_{3,8} - ib_{4,7} + b_{4,8} \rangle \\
L_{-1,1,1,i} = \langle ib_{1,5} - b_{1,6} - ib_{2,5} - b_{2,6} - ib_{3,7} + b_{3,8} + ib_{4,7} + b_{4,8} \rangle \\
L_{1,-1,-1,-i} = \langle -ib_{1,7} - b_{1,8} - ib_{2,7} + b_{2,8} - ib_{3,5} - b_{3,6} - ib_{4,5} + b_{4,6} \rangle \\
L_{1,-1,-1,i} = \langle ib_{1,7} - b_{1,8} + ib_{2,7} + b_{2,8} + ib_{3,5} - b_{3,6} + ib_{4,5} + b_{4,6} \rangle \\
L_{1,-1,1,-i} = \langle ib_{1,7} + b_{1,8} - ib_{2,7} + b_{2,8} + ib_{3,5} + b_{3,6} - ib_{4,5} + b_{4,6} \rangle \\
L_{1,-1,1,i} = \langle -ib_{1,7} + b_{1,8} + ib_{2,7} + b_{2,8} - ib_{3,5} + b_{3,6} + ib_{4,5} + b_{4,6} \rangle \\
L_{1,1,-1,-i} = \langle -ib_{1,5} - b_{1,6} - ib_{2,5} + b_{2,6} - ib_{3,7} - b_{3,8} - ib_{4,7} + b_{4,8} \rangle \\
L_{1,1,-1,i} = \langle ib_{1,5} - b_{1,6} + ib_{2,5} + b_{2,6} + ib_{3,7} - b_{3,8} + ib_{4,7} + b_{4,8} \rangle \\
L_{1,1,1,-i} = \langle ib_{1,5} + b_{1,6} - ib_{2,5} + b_{2,6} + ib_{3,7} + b_{3,8} - ib_{4,7} + b_{4,8} \rangle \\
L_{1,1,1,i} = \langle -ib_{1,5} + b_{1,6} + ib_{2,5} + b_{2,6} - ib_{3,7} + b_{3,8} + ib_{4,7} + b_{4,8} \rangle
\end{array}$$

$$\begin{array}{ll}
L_{-1,-1,-1,-1} = \langle b_{5,8} + b_{6,7} \rangle & L_{-1,-1,-1,1} = \langle -b_{1,3} + b_{2,4} \rangle \\
L_{-1,1,-1,-1} = \langle -b_{1,2} + b_{3,4} \rangle & L_{-1,1,-1,1} = \langle -b_{5,6} + b_{7,8} \rangle \\
L_{1,-1,-1,1} = \langle -b_{5,8} + b_{6,7} \rangle & L_{1,1,-1,1} = \langle b_{5,6} + b_{7,8} \rangle \\
L_{1,1,-1,-1} = \langle b_{1,2} + b_{3,4} \rangle & L_{-1,-1,1,-1} = \langle b_{1,4} + b_{2,3},\ -b_{5,7} + b_{6,8} \rangle \\
L_{1,-1,-1,-1} = \langle -b_{1,4} + b_{2,3} \rangle & L_{-1,-1,1,1} = \langle b_{1,3} + b_{2,4},\ b_{5,7} + b_{6,8} \rangle
\end{array}$$

Grading over $\mathbb{Z}^3 \times \mathbb{Z}_2$: $Q_9 = \langle\{p(u), F_8, r(v), q(w) \mid u, v, w \in \mathbb{C}^*\}\rangle \cong (\mathbb{C}^*)^3 \times \mathbb{Z}_2$ is an abelian diagonalizable subgroup of $\operatorname{Aut} L$, concretely is the MAD-group $T^{(0)}_{6,2}$, which produces a grading of type $(25, 0, 1)$:

$$\begin{array}{ll}
L_{\frac{1}{2},1,1,\frac{1}{3}} = \langle -b_{5,7} + ib_{5,8} + ib_{6,7} + b_{6,8}\rangle & L_{\frac{1}{2},1,\frac{1}{5},1} = \langle -b_{3,7} + ib_{3,8} + ib_{4,7} + b_{4,8}\rangle \\
L_{\frac{1}{2},1,1,3} = \langle b_{5,7} - ib_{5,8} + ib_{6,7} + b_{6,8}\rangle & L_{\frac{1}{2},1,5,1} = \langle b_{3,7} - ib_{3,8} + ib_{4,7} + b_{4,8}\rangle \\
L_{1,1,\frac{1}{5},\frac{1}{3}} = \langle -b_{3,5} + ib_{3,6} + ib_{4,5} + b_{4,6}\rangle & L_{1,1,5,\frac{1}{3}} = \langle b_{3,5} - ib_{3,6} + ib_{4,5} + b_{4,6}\rangle \\
L_{1,1,\frac{1}{5},3} = \langle b_{3,5} + ib_{3,6} - ib_{4,5} + b_{4,6}\rangle & L_{1,1,5,3} = \langle -b_{3,5} - ib_{3,6} - ib_{4,5} + b_{4,6}\rangle \\
L_{2,1,\frac{1}{5},1} = \langle b_{3,7} + ib_{3,8} - ib_{4,7} + b_{4,8}\rangle & L_{2,1,1,\frac{1}{3}} = \langle b_{5,7} + ib_{5,8} - ib_{6,7} + b_{6,8}\rangle \\
L_{2,1,1,3} = \langle -b_{5,7} - ib_{5,8} - ib_{6,7} + b_{6,8}\rangle & L_{2,1,5,1} = \langle -b_{3,7} - ib_{3,8} - ib_{4,7} + b_{4,8}\rangle
\end{array}$$

$$\begin{array}{lll}
L_{\frac{1}{2},-1,1,1} = \langle ib_{1,7} + b_{1,8}\rangle & L_{\frac{1}{2},1,1,1} = \langle ib_{2,7} + b_{2,8}\rangle & L_{1,-1,\frac{1}{5},1} = \langle ib_{1,3} + b_{1,4}\rangle \\
L_{1,-1,1,\frac{1}{3}} = \langle ib_{1,5} + b_{1,6}\rangle & L_{1,-1,1,1} = \langle b_{1,2}\rangle & L_{1,-1,1,3} = \langle -ib_{1,5} + b_{1,6}\rangle \\
L_{1,-1,5,1} = \langle -ib_{1,3} + b_{1,4}\rangle & L_{1,1,\frac{1}{5},1} = \langle ib_{2,3} + b_{2,4}\rangle & L_{1,1,1,\frac{1}{3}} = \langle ib_{2,5} + b_{2,6}\rangle \\
L_{1,1,1,1} = \langle b_{3,4},\, b_{5,6},\, b_{7,8}\rangle & L_{1,1,1,3} = \langle -ib_{2,5} + b_{2,6}\rangle & L_{1,1,5,1} = \langle -ib_{2,3} + b_{2,4}\rangle \\
L_{2,-1,1,1} = \langle -ib_{1,7} + b_{1,8}\rangle & L_{2,1,1,1} = \langle -ib_{2,7} + b_{2,8}\rangle &
\end{array}$$

Grading over $\mathbb{Z}^4$: $Q_{10} = \langle\{\operatorname{Ad} p(u), \operatorname{Ad} s(v), \operatorname{Ad} r(w), \operatorname{Ad} q(z) \mid u, v, w, z \in \mathbb{C}^*\}\rangle \cong (\mathbb{C}^*)^4$ is an abelian diagonalizable subgroup of $\operatorname{Aut} L$ (corresponding to $T^{(0)}_{8,0}$). The simultaneous diagonalization of L relative to the quasitorus Q_{10} is of type $(24, 0, 0, 1)$. Of course it is the *Cartan grading*, in other words, the decomposition in root spaces relative to the Cartan subalgebra $L_{1,1,1,1} = \langle b_{1,2},\, b_{3,4},\, b_{5,6},\, b_{7,8}\rangle$:

$$\begin{array}{ll}
L_{\frac{1}{2},\frac{1}{7},1,1} = \langle -b_{1,7} + ib_{1,8} + ib_{2,7} + b_{2,8}\rangle & L_{\frac{1}{2},1,\frac{1}{5},1} = \langle -b_{3,7} + ib_{3,8} + ib_{4,7} + b_{4,8}\rangle \\
L_{\frac{1}{2},1,1,\frac{1}{3}} = \langle -b_{5,7} + ib_{5,8} + ib_{6,7} + b_{6,8}\rangle & L_{\frac{1}{2},1,1,3} = \langle b_{5,7} - ib_{5,8} + ib_{6,7} + b_{6,8}\rangle \\
L_{\frac{1}{2},1,5,1} = \langle b_{3,7} - ib_{3,8} + ib_{4,7} + b_{4,8}\rangle & L_{\frac{1}{2},7,1,1} = \langle b_{1,7} - ib_{1,8} + ib_{2,7} + b_{2,8}\rangle \\
L_{1,\frac{1}{7},\frac{1}{5},1} = \langle -b_{1,3} + ib_{1,4} + ib_{2,3} + b_{2,4}\rangle & L_{1,\frac{1}{7},1,\frac{1}{3}} = \langle -b_{1,5} + ib_{1,6} + ib_{2,5} + b_{2,6}\rangle \\
L_{1,\frac{1}{7},5,1} = \langle b_{1,3} + ib_{1,4} - ib_{2,3} + b_{2,4}\rangle & L_{1,\frac{1}{7},1,3} = \langle b_{1,5} + ib_{1,6} - ib_{2,5} + b_{2,6}\rangle \\
L_{1,1,\frac{1}{5},\frac{1}{3}} = \langle -b_{3,5} + ib_{3,6} + ib_{4,5} + b_{4,6}\rangle & L_{1,1,\frac{1}{5},3} = \langle b_{3,5} + ib_{3,6} - ib_{4,5} + b_{4,6}\rangle \\
L_{1,1,5,\frac{1}{3}} = \langle b_{3,5} - ib_{3,6} + ib_{4,5} + b_{4,6}\rangle & L_{1,1,5,3} = \langle -b_{3,5} - ib_{3,6} - ib_{4,5} + b_{4,6}\rangle \\
L_{1,7,\frac{1}{5},1} = \langle b_{1,3} - ib_{1,4} + ib_{2,3} + b_{2,4}\rangle & L_{1,7,1,\frac{1}{3}} = \langle b_{1,5} - ib_{1,6} + ib_{2,5} + b_{2,6}\rangle \\
L_{1,7,5,1} = \langle -b_{1,3} - ib_{1,4} - ib_{2,3} + b_{2,4}\rangle & L_{1,7,1,3} = \langle -b_{1,5} - ib_{1,6} - ib_{2,5} + b_{2,6}\rangle \\
L_{2,\frac{1}{7},1,1} = \langle b_{1,7} + ib_{1,8} - ib_{2,7} + b_{2,8}\rangle & L_{2,1,\frac{1}{5},1} = \langle b_{3,7} + ib_{3,8} - ib_{4,7} + b_{4,8}\rangle \\
L_{2,1,1,\frac{1}{3}} = \langle b_{5,7} + ib_{5,8} - ib_{6,7} + b_{6,8}\rangle & L_{2,1,1,3} = \langle -b_{5,7} - ib_{5,8} - ib_{6,7} + b_{6,8}\rangle \\
L_{2,1,5,1} = \langle -b_{3,7} - ib_{3,8} - ib_{4,7} + b_{4,8}\rangle & L_{2,7,1,1} = \langle -b_{1,7} - ib_{1,8} - ib_{2,7} + b_{2,8}\rangle
\end{array}$$

Grading over $\mathbb{Z}_2^6$: $Q_{11} = \langle\{G_8, G_{10}, G_{13}, G_3, G_7, G_{14}\}\rangle \cong \mathbb{Z}_2^6$ is also a MAD of $\operatorname{Aut} L$, corresponding to $T^{(3)}_{0,1}$. The induced grading is of type (28):

$$\begin{array}{ll}
L_{-1,-1,-1,-1,-1,-1} = \langle b_{1,8} - b_{2,7} - b_{3,6} + b_{4,5}\rangle & L_{-1,-1,-1,-1,1,1} = \langle b_{1,7} - b_{2,8} - b_{3,5} + b_{4,6}\rangle \\
L_{-1,-1,-1,1,-1,1} = \langle -b_{1,8} - b_{2,7} + b_{3,6} + b_{4,5}\rangle & L_{-1,-1,-1,1,1,1} = \langle -b_{1,7} - b_{2,8} + b_{3,5} + b_{4,6}\rangle \\
L_{-1,-1,1,-1,1,-1} = \langle b_{1,3} - b_{2,4} - b_{5,7} + b_{6,8}\rangle & L_{-1,-1,1,-1,1,1} = \langle -b_{1,3} + b_{2,4} - b_{5,7} + b_{6,8}\rangle \\
L_{-1,-1,1,1,-1,-1} = \langle -b_{1,4} - b_{2,3} + b_{5,8} + b_{6,7}\rangle & L_{-1,-1,1,1,-1,1} = \langle b_{1,4} + b_{2,3} + b_{5,8} + b_{6,7}\rangle \\
L_{-1,-1,1,1,1,-1} = \langle -b_{1,3} - b_{2,4} + b_{5,7} + b_{6,8}\rangle & L_{-1,-1,1,1,1,1} = \langle b_{1,3} + b_{2,4} + b_{5,7} + b_{6,8}\rangle \\
L_{-1,1,-1,-1,-1,1} = \langle b_{1,6} - b_{2,5} - b_{3,8} + b_{4,7}\rangle & L_{-1,1,-1,-1,1,-1} = \langle b_{1,5} - b_{2,6} - b_{3,7} + b_{4,8}\rangle \\
L_{-1,1,-1,1,-1,-1} = \langle -b_{1,6} - b_{2,5} + b_{3,8} + b_{4,7}\rangle & L_{-1,1,-1,1,1,-1} = \langle -b_{1,5} - b_{2,6} + b_{3,7} + b_{4,8}\rangle \\
L_{-1,1,1,-1,-1,-1} = \langle b_{1,2} - b_{3,4} - b_{5,6} + b_{7,8}\rangle & L_{-1,1,1,-1,-1,1} = \langle -b_{1,2} + b_{3,4} - b_{5,6} + b_{7,8}\rangle \\
L_{1,-1,-1,-1,-1,1} = \langle -b_{1,8} + b_{2,7} - b_{3,6} + b_{4,5}\rangle & L_{1,-1,-1,-1,1,-1} = \langle -b_{1,7} + b_{2,8} - b_{3,5} + b_{4,6}\rangle \\
L_{1,-1,-1,1,-1,-1} = \langle b_{1,8} + b_{2,7} + b_{3,6} + b_{4,5}\rangle & L_{1,-1,-1,1,1,-1} = \langle b_{1,7} + b_{2,8} + b_{3,5} + b_{4,6}\rangle \\
L_{1,-1,1,-1,-1,-1} = \langle b_{1,4} - b_{2,3} - b_{5,8} + b_{6,7}\rangle & L_{1,-1,1,-1,-1,1} = \langle -b_{1,4} + b_{2,3} - b_{5,8} + b_{6,7}\rangle \\
L_{1,1,-1,-1,-1,1} = \langle -b_{1,6} + b_{2,5} - b_{3,8} + b_{4,7}\rangle & L_{1,1,-1,-1,1,-1} = \langle -b_{1,5} + b_{2,6} - b_{3,7} + b_{4,8}\rangle \\
L_{1,1,-1,1,-1,-1} = \langle b_{1,6} + b_{2,5} + b_{3,8} + b_{4,7}\rangle & L_{1,1,-1,1,1,-1} = \langle b_{1,5} + b_{2,6} + b_{3,7} + b_{4,8}\rangle \\
L_{1,1,1,-1,-1,-1} = \langle -b_{1,2} - b_{3,4} + b_{5,6} + b_{7,8}\rangle & L_{1,1,1,-1,-1,1} = \langle b_{1,2} + b_{3,4} + b_{5,6} + b_{7,8}\rangle
\end{array}$$

In order to describe the gradings involving a copy of $\mathbb{Z}_3$, we introduce new automorphisms. Fix $\mathcal{B} = \{b_1, \dots, b_{28}\}$ a basis of L of root vectors (take the basis produced by Q_{10}, suitably ordered, to express the remaining three gradings also in terms of b_{ij}), and let $\varphi_{i,j}(b_k) = \delta_{k,j} b_i$ for $i, j, k = 1 \dots 28$. Define

$$
\begin{aligned}
H_1 = {}& \varphi_{2,4} + \varphi_{3,1} + \varphi_{3,2} + \varphi_{5,27} + \varphi_{6,8} + \varphi_{7,9} + \varphi_{8,25} + \varphi_{11,12} + \varphi_{12,26} + \varphi_{13,18} + \varphi_{14,23} \\
& + \varphi_{17,15} + \varphi_{18,20} + \varphi_{19,21} + \varphi_{20,13} + \varphi_{23,24} + \varphi_{24,14} + \varphi_{25,6} + \varphi_{26,11} - \varphi_{1,1} - \varphi_{1,2} - \varphi_{1,3} \\
& - \varphi_{1,4} - \varphi_{4,2} - \varphi_{4,4} - \varphi_{9,22} - \varphi_{10,19} - \varphi_{15,28} - \varphi_{16,5} - \varphi_{21,10} - \varphi_{22,7} - \varphi_{27,16} - \varphi_{28,17} \\
H_2 = {}& \varphi_{2,2} + \varphi_{2,3} + \varphi_{2,4} + \varphi_{4,1} + \varphi_{5,26} + \varphi_{6,16} + \varphi_{7,19} + \varphi_{8,5} + \varphi_{10,22} + \varphi_{11,13} + \varphi_{12,18} + \varphi_{14,8} \\
& + \varphi_{15,23} + \varphi_{16,12} + \varphi_{17,14} + \varphi_{18,28} + \varphi_{19,7} + \varphi_{20,17} + \varphi_{22,10} + \varphi_{23,25} + \varphi_{24,6} + \varphi_{26,20} \\
& + \varphi_{27,11} + \varphi_{28,24} - \varphi_{1,1} - \varphi_{1,3} - \varphi_{1,4} - \varphi_{3,3} - \varphi_{9,21} - \varphi_{13,15} - \varphi_{21,9} - \varphi_{25,27} - 2\varphi_{1,2} \\
t_{x,y,z,u} := {}& \operatorname{diag}\{x, y, z, u, xy, xyz, yz, uxy, uy, uxy^2z, uxyz, uyz, \frac{1}{x}, \frac{1}{y}, \frac{1}{z}, \frac{1}{u}, \frac{1}{xy}, \frac{1}{xyz}, \frac{1}{yz}, \frac{1}{uxy}, \\
& \frac{1}{uy}, \frac{1}{uxy^2z}, \frac{1}{uxyz}, \frac{1}{uyz}\}
\end{aligned}
$$

Grading over $\mathbb{Z}_3^3$: If we denote $\omega = -(-1)^{\frac{1}{3}}$, $Q_{12} = \langle\{t_{1,\omega^2,\omega^2,\omega^2}, t_{\omega^2,1,\omega^2,1}, H_1\}\rangle \cong \mathbb{Z}_3^3$ is an abelian diagonalizable subgroup of $\operatorname{Aut} L$ inducing a grading of type $(24, 2)$:

$$
\begin{aligned}
L_{1,1,\omega} &= \langle -\omega b_1 - \omega b_2 + b_3,\ (1+\omega) b_2 + b_4 \rangle \\
L_{1,1,\omega^2} &= \langle -\omega^2 b_1 - \omega^2 b_2 + b_3,\ (1+\omega^2) b_2 + b_4 \rangle
\end{aligned}
$$

$$
\begin{aligned}
L_{1,\omega,1} &= \langle b_{13} + b_{18} + b_{20} \rangle & L_{1,\omega,\omega} &= \langle \omega^2 b_{13} + \omega b_{18} + b_{20} \rangle \\
L_{1,\omega,\omega^2} &= \langle \omega b_{13} + \omega^2 b_{18} + b_{20} \rangle & L_{1,\omega^2,1} &= \langle b_6 + b_8 + b_{25} \rangle \\
L_{1,\omega^2,\omega} &= \langle \omega^2 b_6 + \omega b_8 + b_{25} \rangle & L_{1,\omega^2,\omega^2} &= \langle \omega b_6 + \omega^2 b_8 + b_{25} \rangle \\
L_{\omega,1,1} &= \langle -b_{15} - b_{17} + b_{28} \rangle & L_{\omega,1,\omega} &= \langle -\omega b_{15} - \omega^2 b_{17} + b_{28} \rangle \\
L_{\omega,1,\omega^2} &= \langle -\omega^2 b_{15} - \omega b_{17} + b_{28} \rangle & L_{\omega,\omega,1} &= \langle -b_{10} + b_{19} + b_{21} \rangle \\
L_{\omega,\omega,\omega} &= \langle -\omega^2 b_{10} + \omega b_{19} + b_{21} \rangle & L_{\omega,\omega,\omega^2} &= \langle -\omega b_{10} + \omega^2 b_{19} + b_{21} \rangle \\
L_{\omega,\omega^2,1} &= \langle b_{14} + b_{23} + b_{24} \rangle & L_{\omega,\omega^2,\omega} &= \langle \omega^2 b_{14} + \omega b_{23} + b_{24} \rangle \\
L_{\omega,\omega^2,\omega^2} &= \langle \omega b_{14} + \omega^2 b_{23} + b_{24} \rangle & L_{\omega^2,1,1} &= \langle b_5 - b_{16} + b_{27} \rangle \\
L_{\omega^2,1,\omega} &= \langle \omega b_5 - \omega^2 b_{16} + b_{27} \rangle & L_{\omega^2,1,\omega^2} &= \langle \omega^2 b_5 - \omega b_{16} + b_{27} \rangle \\
L_{\omega^2,\omega,1} &= \langle b_{11} + b_{12} + b_{26} \rangle & L_{\omega^2,\omega,\omega} &= \langle \omega^2 b_{11} + \omega b_{12} + b_{26} \rangle \\
L_{\omega^2,\omega,\omega^2} &= \langle \omega b_{11} + \omega^2 b_{12} + b_{26} \rangle & L_{\omega^2,\omega^2,1} &= \langle -b_7 - b_9 + b_{22} \rangle \\
L_{\omega^2,\omega^2,\omega} &= \langle -\omega^2 b_7 - \omega b_9 + b_{22} \rangle & L_{\omega^2,\omega^2,\omega^2} &= \langle -\omega b_7 - \omega^2 b_9 + b_{22} \rangle
\end{aligned}
$$

Grading over $\mathbb{Z}^2 \times \mathbb{Z}_3$: $Q_{13} = \langle\{t_{u,1,u,\frac{1}{u}}, t_{1,v,\frac{1}{v^2},1}, H_2^2 \mid u, v \in \mathbb{C}^*\}\rangle \cong (\mathbb{C}^*)^2 \times \mathbb{Z}_3$ is an abelian diagonalizable subgroup of $\operatorname{Aut} L$, which produces a grading of type $(26, 1)$:

$$
\begin{aligned}
L_{\frac{1}{2},1,1} &= \langle b_8 + b_{17} + b_{26} \rangle & L_{\frac{1}{2},1,\omega} &= \langle \omega b_8 + \omega^2 b_{17} + b_{26} \rangle \\
L_{\frac{1}{2},1,\omega^2} &= \langle \omega^2 b_8 + \omega b_{17} + b_{26} \rangle & L_{\frac{1}{2},3,1} &= \langle b_{13} - b_{23} + b_{27} \rangle \\
L_{\frac{1}{2},3,\omega} &= \langle \omega^2 b_{13} - \omega b_{23} + b_{27} \rangle & L_{\frac{1}{2},3,\omega^2} &= \langle \omega b_{13} - \omega^2 b_{23} + b_{27} \rangle \\
L_{1,\frac{1}{3},1} &= \langle b_{16} + b_{18} + b_{24} \rangle & L_{1,\frac{1}{3},\omega} &= \langle \omega^2 b_{16} + \omega b_{18} + b_{24} \rangle \\
L_{1,1,\omega} &= \langle \tfrac{1+\omega^2}{1-\omega} b_1 + \tfrac{2}{1-\omega} b_2 + \tfrac{1}{1-\omega} b_3 + b_4 \rangle & L_{1,1,\omega^2} &= \langle \tfrac{1}{1-\omega} b_1 + \tfrac{2}{1-\omega^2} b_2 + \tfrac{1}{1-\omega^2} b_3 + b_4 \rangle \\
L_{1,3,1} &= \langle b_6 + b_{12} + b_{28} \rangle & L_{1,3,\omega} &= \langle \omega^2 b_6 + \omega b_{12} + b_{28} \rangle \\
L_{1,3,\omega^2} &= \langle \omega b_6 + \omega^2 b_{12} + b_{28} \rangle & L_{2,\frac{1}{3},1} &= \langle -b_{11} + b_{15} + b_{25} \rangle \\
L_{2,\frac{1}{3},\omega} &= \langle -\omega^2 b_{11} + \omega b_{15} + b_{25} \rangle & L_{2,\frac{1}{3},\omega^2} &= \langle -\omega b_{11} + \omega^2 b_{15} + b_{25} \rangle \\
L_{2,1,1} &= \langle b_5 + b_{14} + b_{20} \rangle & L_{2,1,\omega} &= \langle \omega b_5 + \omega^2 b_{14} + b_{20} \rangle \\
L_{1,\frac{1}{3},\omega^2} &= \langle \omega b_{16} + \omega^2 b_{18} + b_{24} \rangle & L_{2,1,\omega^2} &= \langle \omega^2 b_5 + \omega b_{14} + b_{20} \rangle
\end{aligned}
$$

$$
\begin{array}{llll}
L_{1,1,1} = \langle b_1 + b_2,\ b_3 \rangle & L_{\frac{1}{4},3,1} = \langle b_{22} \rangle & \text{Ł}_{\frac{1}{2},\frac{1}{3},1} = \langle b_{21} \rangle & \\
L_{\frac{1}{2},9,1} = \langle b_{19} \rangle & L_{2,\frac{1}{9},1} = \langle b_7 \rangle & L_{2,3,1} = \langle b_9 \rangle & L_{4,\frac{1}{3},1} = \langle b_{10} \rangle
\end{array}
$$

Grading over $\mathbb{Z}_2^3 \times \mathbb{Z}_3$: $Q_{14} = \langle \{t_{1,-1,1,1}, t_{-1,-1,-1,-1}, H_2\}\rangle \cong \mathbb{Z}_2^2 \times \mathbb{Z}_6$ is an abelian diagonalizable subgroup of $\operatorname{Aut} L$, producing a grading of type $(14, 7)$:

$$
\begin{aligned}
&L_{-1,-1,-1} = \langle b_6 + b_{12} - b_{16} - b_{18} - b_{24} + b_{28},\ -b_{10} + b_{22}\rangle\\
&L_{-1,-1,1} = \langle b_6 + b_{12} + b_{16} + b_{18} + b_{24} + b_{28},\ b_{10} + b_{22}\rangle\\
&L_{-1,1,-1} = \langle -b_{11} + b_{13} + b_{15} - b_{23} + b_{25} + b_{27},\ b_9 + b_{21}\rangle\\
&L_{-1,1,1} = \langle b_{11} + b_{13} - b_{15} - b_{23} - b_{25} + b_{27},\ -b_9 + b_{21}\rangle\\
&L_{1,-1,-1} = \langle -b_5 + b_8 - b_{14} + b_{17} - b_{20} + b_{26},\ -b_7 + b_{19}\rangle\\
&L_{1,-1,1} = \langle b_5 + b_8 + b_{14} + b_{17} + b_{20} + b_{26},\ b_7 + b_{19}\rangle\\
&L_{1,1,-1} = \langle b_3,\ b_1 + b_2\rangle\\
&L_{-1,-1,\omega} = \langle \omega b_6 + \omega^2 b_{12} + b_{16} + \omega b_{18} + \omega^2 b_{24} + b_{28}\rangle\\
&L_{-1,-1,\ -\omega} = \langle \omega b_6 + \omega^2 b_{12} - b_{16} - \omega b_{18} - \omega^2 b_{24} + b_{28}\rangle\\
&L_{-1,-1,-\omega^2} = \langle \omega^2 b_6 + \omega b_{12} - b_{16} - \omega^2 b_{18} - \omega b_{24} + b_{28}\rangle\\
&L_{-1,-1,\omega^2} = \langle \omega^2 b_6 + \omega b_{12} + b_{16} + \omega^2 b_{18} + \omega b_{24} + b_{28}\rangle\\
&L_{-1,1,\omega} = \langle \omega^2 b_{11} + \omega b_{13} - b_{15} - \omega^2 b_{23} - \omega b_{25} + b_{27}\rangle\\
&L_{-1,1,-\omega} = \langle -\omega^2 b_{11} + \omega b_{13} + b_{15} - \omega^2 b_{23} + \omega b_{25} + b_{27}\rangle\\
&L_{-1,1,-\omega^2} = \langle -\omega b_{11} + \omega^2 b_{13} + b_{15} - \omega b_{23} + \omega^2 b_{25} + b_{27}\rangle\\
&L_{-1,1,\omega^2} = \langle \omega b_{11} + \omega^2 b_{13} - b_{15} - \omega b_{23} - \omega^2 b_{25} + b_{27}\rangle\\
&L_{1,-1,\omega} = \langle \omega b_5 + \omega^2 b_8 + b_{14} + \omega b_{17} + \omega^2 b_{20} + b_{26}\rangle\\
&L_{1,-1,-\omega} = \langle -\omega b_5 + \omega^2 b_8 - b_{14} + \omega b_{17} - \omega^2 b_{20} + b_{26}\rangle\\
&L_{1,-1,-\omega^2} = \langle -\omega^2 b_5 + \omega b_8 - b_{14} + \omega^2 b_{17} - \omega b_{20} + b_{26}\rangle\\
&L_{1,-1,\omega^2} = \langle \omega^2 b_5 + \omega b_8 + b_{14} + \omega^2 b_{17} + \omega b_{20} + b_{26}\rangle\\
&L_{1,1,-\omega} = \langle \tfrac{-\omega-\omega^2}{1-\omega} b_1 - \tfrac{2\omega}{1-\omega} b_2 - \tfrac{\omega}{1-\omega} b_3 + b_4\rangle\\
&L_{1,1,-\omega^2} = \langle \tfrac{-\omega}{1-\omega} b_1 + \tfrac{2\omega^2}{-1+\omega^2} b_2 + \tfrac{\omega^2}{-1+\omega^2} b_3 + b_4\rangle
\end{aligned}
$$

References

[1] Y. A. Bahturin, I. P. Shestakov and M. V. Zaicev: *Journal of Algebra* **283**, 849–868 (2005).

[2] C. Draper and C. Martín: *Linear Algebra and its Applications* **418**, 85–111 (2006).

[3] C. Draper and C. Martín: *Gradings on the Albert algebra and on* $\mathfrak{f}_4$. Preprint arXiv:math-ra/0703840v1.

[4] C. Draper and A. Viruel: *Gradings on* $\mathfrak{o}_4$. Preprint.

[5] M. Havlcek, J. Patera and E. Pelantová: *Linear Algebra and its Applications* **277**, 97–125 (1998).

[6] P. Jordan: *Nachr. Ges. Wiss. Göttingen* (1933), 209-217.

[7] A.U. Klimyk: *Czech. J. Phys.* **51** 331 (2001).

[8] J. Patera and H. Zassenhaus: *Linear Algebra and its Applications* **112** 87–159 (1989).

[9] J. Patera, E. Pelantová and M. Svobodová: *J. Math. Phys.* **42**, 3839-3853 (2001).

[10] J. Patera, E. Pelantová and M. Svobodová: *J. Math. Phys.* **43**, 6353-6378 (2002).

[11] J. Patera, E. Pelantová and M. Svobodová: *J. Math. Phys.* **45**, 2188-2198 (2004).

[12] J. Patera, E. Pelantová and M. Svobodová: *Fine gradings of the real forms of* $sl(4,\mathbb{C})$, $sp(4,\mathbb{C})$, *and* $o(4,\mathbb{C})$. Preprint arXiv:math-ph/0608032v1.